



## Neural Mechanisms of Hyperalgesia After Tissue Injury

Richard A. Meyer, Matthias Ringkamp, James N. Campbell, and Srinivasa N. Raja

**F**or more than 25 years, investigators from APL and the Johns Hopkins School of Medicine have collaborated on research aimed at understanding the neural mechanisms of pain sensation. This research is based on correlating results from studies of pain sensations in humans with results from studies of neural activity in anesthetized animals. One aspect of pain that has clinical importance is hyperalgesia—the enhanced pain to stimuli applied to the skin that develops after tissue injury and in certain diseases. We review here the neural mechanisms of hyperalgesia. Primary hyperalgesia, which develops at the site of tissue injury, is associated with an increased sensitivity of the peripheral nerve fibers involved in pain. Secondary hyperalgesia, which develops in uninjured tissue surrounding the site of injury, exhibits symptoms similar to those seen in chronic pain patients and is caused by an enhanced neural responsiveness in the central nervous system.


### INTRODUCTION

Pain serves as a warning system to protect an organism from injury and is one of the main reasons that patients go to physicians. In most cases, treatment of the underlying injury or disease ameliorates the pain, but for some, the pain persists well after the disease or injury has been cured. This persistent pain is called chronic pain and does not serve a useful biological function.

APL has been interested in a form of chronic pain called neuropathic pain. In this disease, pathology in the nervous system leads to chronic pain. The pathology can be caused by a disease (e.g., diabetes, shingles, AIDS) or trauma to the nerve (e.g., from a bullet or knife wound). Patients with neuropathic pain complain of ongoing pain as well as enhanced pain sensitivity to stimuli applied to their skin. For example, lightly

touching the skin can be extremely painful in some patients with neuropathic pain. We call this increased pain sensitivity hyperalgesia (from the Greek, *hyper* = above, *algos* = pain).

As an example, an elderly woman in otherwise good health experiences the sudden onset of a localized region of shooting pain in the left side of the chest, starting from the midline in the back and radiating toward the front. Two days later she notices an inch-wide band-like red rash with scattered small blisters in the same region of the back and chest (Fig. 1a). She consults her physician, who diagnoses her condition as an acute attack of shingles (herpes zoster) and tells her that her symptoms should subside in a few days and that the skin lesions will heal in 2–3 weeks. Six months later, the patient continues to have pain and hypersensitivity in the



**Figure 1.** Pain and hyperalgesia in a patient with postherpetic neuralgia. (a) This elderly patient reported pain with skin eruptions along a narrow band on her back and chest. She was diagnosed to have herpes zoster (shingles). (b) Six months later, after the lesions had healed, her pain persisted. She was diagnosed to have postherpetic neuralgia, a form of neuropathic pain. She complained of ongoing pain, pain to a light brush (hyperalgesia) within the zone indicated by the red border, and pain to punctate stimuli within the zone indicated by the blue border.

chest; the area of pain now extends to involve a large area on the back (Fig. 1b). Even brushing the skin with a cotton swab is painful, as are clothes or sheets rubbing against her skin in the affected region. Thus, a localized region of pain associated with an episode of shingles has resulted in a widespread area of intense pain and hypersensitivity such that even touch and pressure are painful. This phenomenon of hyperalgesia characterizes many of the chronic pain states associated with disease or injury that affect the peripheral or central nervous system.


Hyperalgesia also develops after injury to the skin. Most readers are familiar with the enhanced pain that develops after a sunburn; just the rubbing of fabric against the skin and gentle warming from a shower cause discomfort. The hyperalgesia that develops after skin injury is similar in many aspects to the hyperalgesia that develops in neuropathic pain. Studies of hyperalgesia after skin injury therefore provide a human surrogate for studies of hyperalgesia in neuropathic pain. In this article, we briefly review the mechanism of hyperalgesia that develops after an injury to the skin, focusing mainly on changes which occur in the peripheral nervous system that may account for the hyperalgesia.

## TWO FORMS OF HYPERALGESIA

When a natural stimulus (e.g., heat) is applied to the skin, there is a range of stimulus intensities over which the stimulus is not painful. As the stimulus intensity is increased, however, a stimulus level


is eventually reached where the stimulus becomes painful. Above this level, pain increases with stimulus intensity. A hypothetical stimulus-response function is shown in Fig. 2. When the skin is injured and hyperalgesia develops, this stimulus-response function is shifted to the left, resulting in a lowering of the pain threshold and an increase in pain to suprathreshold stimuli. We seek to determine what in the nervous system can account for this leftward shift in the stimulus-response function.

Injury to the skin (e.g., from a burn) can lead to the development of two forms of hyperalgesia (Fig. 3). Primary hyperalgesia develops at the site of the injury. Secondary



**Figure 2.** Hyperalgesia. Magnitude of pain is plotted as a function of stimulus intensity. In normal skin, pain increases with stimulus intensity above the pain threshold. For hyperalgesic skin, the stimulus-response function is shifted to the left; there is a lower threshold for producing pain and an increased response to suprathreshold stimuli.

	Primary	Secondary
Mechanical hyperalgesia	Yes	Yes
Heat hyperalgesia	Yes	No



**Figure 3.** Primary and secondary hyperalgesia differ. After a tissue injury, primary hyperalgesia develops at the site of injury and secondary hyperalgesia develops in uninjured tissue surrounding the site of injury.

hyperalgesia develops in the uninjured tissue that surrounds the injury. The characteristics of primary and secondary hyperalgesia differ. In the area of primary hyperalgesia, there is hyperalgesia to both mechanical and heat stimuli. In contrast, in the area of secondary hyperalgesia, there is hyperalgesia to mechanical stimuli but not to heat stimuli.<sup>1,2</sup> This dichotomy suggests that the neural mechanisms of primary and secondary hyperalgesia differ.

### Primary Hyperalgesia

In one of our early studies to investigate the mechanisms of primary hyperalgesia, we applied a controlled burn injury to the hand.<sup>3</sup> The injury was produced by a 53°C heat stimulus applied for 30 s using a laser thermal stimulator system developed at APL (see the boxed insert below). This stimulus produced intense pain and resulted in a blister several hours later in about half of the subjects (Fig. 4).

#### LASER THERMAL STIMULATOR


To study pain quantitatively it is necessary to be able to deliver well-controlled noxious stimuli to the skin. Pain can be produced by noxious heat, cold, mechanical, and chemical stimuli. More than 30 years ago, a laser thermal stimulator system was developed at APL to deliver constant temperature stimuli to the skin.<sup>4</sup>

The laser thermal stimulator system consists of a 50-W carbon dioxide laser as the heat source and an IR radiometer to remotely sense skin temperature (Fig. A). The laser operates in a fundamental mode (called TEM<sub>00</sub>), resulting in a beam intensity profile that is a Gaussian of revolution. To achieve a relatively uniform heating profile on the skin, a circular aperture is used to block all but the central 10% of the beam. A germanium lens is used to form an image of this aperture on the skin. The beam diameter on the skin is about 8 mm.



The radiometer consists of a 4-in.-dia. Cassegrain optical system that focuses the target onto an indium antimonide detector system cooled to liquid nitrogen temperature. The optical axis of the radiometer is co-linear with the laser. The beamwidth of the radiometer (6.7-mm dia.) is smaller than the laser spot on the skin. A spectral filter restricts the energy on the detector to a 3- to 5- $\mu\text{m}$  bandwidth. At a 400-Hz bandwidth, the noise equivalent temperature of the system is approximately 0.05°C.

An articulated light guide allows the stimulus to be directed at the target. A two-axis, motorized mirror system at the end of the light guide allows the stimulus to be moved remotely. A co-linear, visible HeNe laser is used for alignment.

This system, shown in Fig. B, is still in active use.



**Figure A.** Block diagram of the APL-developed laser thermal stimulator system. (Adapted from Ref. 4 with permission, © 1976, IEEE.)



**Figure B.** (top) Photograph of the laser thermal stimulator. The flat platform at the top serves as an optical table on which are mounted the carbon dioxide and HeNe lasers and the beam-shaping optics. (bottom) A typical stimulus waveform.




**Figure 4.** A controlled heat injury was used to study primary hyperalgesia. A 53°C heat stimulus was applied to the nonhairy surface of the hand for 30 s. This resulted in a blister several hours later in about half of the subjects.

The laser was also used to test heat sensitivity at the site of the burn before and after the burn injury. Temperatures ranging from 41° to 49°C (3-s duration) were applied to the hand. Subjects used the technique of magnitude estimation (see the boxed insert below) to

#### MEASUREMENTS OF PAIN

Several techniques are available to assess the intensity of pain in humans. A common method is the visual analog scale (VAS), in which verbal descriptors of pain are placed along a scale and subjects mark the scale at a level corresponding to their pain (Fig. A). Subjects use a mouse to move the bar up and down the scale, and ratings are digitized at fixed time points. A method we have used often because it is amenable to correlation with neurophysiological data is the magnitude estimation technique. Here, subjects assign an arbitrary number (e.g., 10) to the intensity of pain associated with a standard stimulus (e.g., 45°C) and rate the intensity of subsequent stimuli relative to this modulus. For example, if the next stimulus was twice as painful as the standard stimulus, subjects would report twice the value of the modulus (i.e., 20 in this example) or half as painful, half the modulus (i.e., 5 in this example). Thus, all ratings are based on a ratio with respect to the modulus. The data in Fig. B were collected from human subjects in response to heat stimuli. The first stimulus was always 45°C. The remaining stimuli ranged from 41° to 49°C and were presented in random order. Subjects used the first stimulus as their modulus and rated the subsequent stimuli relative to this first stimulus. The same stimulus sequence was presented to C-fiber mechano-heat-sensitive nociceptors (CMHs) in deeply anesthetized monkeys (Fig. B). A skin incision was made along the course of the peripheral nerve of interest, and the nerve was dissected from connective tissue. A longitudinal slit was made in the tissue surrounding the nerve trunk, and a small bundle of nerve fibers was cut away from the nerve and rotated onto a dissection platform. Under an operation microscope, the nerve bundle was teased apart with jewelers' forceps to obtain fine filaments with only a few nerve fibers. A small filament was placed on a silver wire electrode, and neural activity was recorded using a low-noise differential amplifier. Neural activity consisted of impulses of electrical signals called action potentials. The skin in the distribution of the monkey nerve under study was squeezed until a region was found where squeezing evoked a response in one of the nerve fibers on the recording electrode. Nylon monofilaments were




**Figure A.** A VAS to measure pain in humans. Pain intensity is rated using a mouse to move the bar along a scale displayed on a computer monitor. (Reprinted from Ref. 5 with permission, © 1995, International Association for the Study of Pain.)

used to determine the area on the skin where the nerve fiber was responsive (the "receptive field"). The nylon monofilaments

were bent at a force that depended on the filament diameter. We used calibrated monofilaments of different diameters to determine the threshold force to achieve a response. Subsequent mechanical, thermal, and chemical stimuli were then applied at or near the receptive field.

Data from the human psychophysical experiments and the monkey neurophysiological experiments were normalized by dividing by the response to the first stimulus. The close correlation in the stimulus-response functions for the human pain ratings and the CMH responses provide strong evidence that heat pain is signaled by activity in these nerve fibers. Note in Fig. B that the normalized response to the 45°C stimulus delivered as part of the random sequence is much less than 1, reflecting the pronounced fatigue that occurs to repeated heat stimulation at short interstimulus intervals (in this case, 30 s).<sup>7</sup>



**Figure B.** C-fiber nociceptors signal heat pain from the human subjects' hands. The magnitude estimation technique was used by subjects to rate the intensity of pain to heat stimuli ranging from 41° to 49°C. The same temperatures were presented to the terminals of C-fiber nociceptors in anesthetized monkeys. The close correlation of these two curves provides evidence that C-fiber nociceptors are responsible for heat pain sensation (from Ref. 6).




rate the intensity of pain to these heat stimuli. Before injury, the pain threshold was around 45°C and pain ratings increased monotonically with stimulus intensity above pain threshold (Fig. 5a). A marked hyperalgesia to heat developed after the burn injury as was evident by a leftward shift in the stimulus-response function. There was a decrease in pain threshold and an increase in pain to suprathreshold stimuli. For example, the 41°C stimulus was not painful before the injury. After the injury, however, the 41°C stimulus was more painful than the 49°C stimulus was before injury.

What accounts for this primary hyperalgesia to heat that develops at the site of injury? Are there changes in the response properties of the nerve fibers that go to the area of injured skin? To address these questions, we recorded the neural activity from single nerve fibers in the peripheral nerve of an anesthetized monkey (see

"Measurements of Pain"). We recorded from two classes of peripheral nerve fibers called A-fibers and C-fibers. This classification is based on the speed at which signals in the nerve fibers propagate from the periphery to the spinal cord; A-fibers are myelinated and propagate at fast conduction velocities (2–50 m/s), whereas C-fibers are unmyelinated and propagate at slow conduction velocities (0.5–2.0 m/s). We were particularly interested in nerve fibers that had receptors in the skin responding to intense, noxious stimuli (i.e., "nociceptors," from the Latin, *noceo* = to injure, hurt). Nociceptors are thought to provide the peripheral signal for pain sensation. We recorded neural activity from nociceptors that responded to both mechanical and heat stimuli: AMHs (A-fiber mechano-heat-sensitive nociceptors) and CMHs (C-fiber mechano-heat-sensitive nociceptors).

The average responses of AMHs and CMHs to heat stimuli before and after the burn injury are shown in Figs. 5b and 5c, respectively. We first focus on the response of the CMHs. Before the injury, as noted previously (Fig. 5c), the threshold for response is around 45°C, and the response increases monotonically with stimulus temperature. The stimulus-response function of the CMHs (Fig. 5c) correlates well with the pain ratings before injury (Fig. 5a) and provides evidence that CMHs



**Figure 5.** Sensitization of peripheral nociceptors accounts for primary hyperalgesia to heat on the hand. (a) Pain ratings to heat stimuli before and after the burn injury. Before injury, the heat threshold was around 45°C. After injury, there was a leftward shift in the stimulus-response function, and marked hyperalgesia to heat developed. (b) Responses of A-fiber mechano-heat-sensitive nociceptors (AMHs) to heat stimuli before and after the burn injury. Before injury, most of the AMHs did not respond to the test heat stimuli. After injury, the AMHs were markedly sensitized to heat. The heat threshold was lowered and the responses to suprathreshold stimuli were enhanced. (c) Responses of C-fiber mechano-heat-sensitive nociceptors (CMHs) to heat stimuli before and after the burn injury. Before injury, the heat threshold was around 45°C and the stimulus-response curve was quite similar to the stimulus-response curve for the human pain ratings. This provides evidence that CMHs signal heat pain from normal skin (see "Measurements of Pain"). However, after injury, the CMHs were desensitized to heat stimuli. The heat threshold increased and the response to suprathreshold stimuli was lower. These results suggest that sensitization of AMHs is responsible for the heat hyperalgesia that develops after a burn to the hand. (Reprinted from Ref. 3 with permission, © 1981, AAAS.)

encoded the pain to heat before injury. After injury, the stimulus-response function of the CMHs is shifted to the right. The threshold is higher, and the response to suprathreshold stimuli is decreased. This is opposite to what would be expected from the observed pain ratings and suggests that CMHs do not provide the neuronal signal for heat hyperalgesia from the injury site.

Now consider the AMHs. Before injury, most AMHs do not respond to heat over the range of temperatures tested (Fig. 5b). However after injury, there is a marked sensitization in the heat response. The threshold for response is lowered and the magnitude of the response to suprathreshold stimuli is increased. These results provide evidence that sensitization of AMHs accounts for the primary hyperalgesia to heat that develops after an injury.


These studies were done on the glabrous (i.e., non-hairy) skin of the hand. Burn injuries to the hairy skin of the arm also lead to hyperalgesia to heat. However, in contrast to glabrous skin, burn injuries to hairy skin cause a sensitization of both CMHs and AMHs. We currently do not know what accounts for this difference. Regardless, in general, primary hyperalgesia is thought to be due to an enhanced responsiveness or sensitization of the nociceptors that signal pain.

### Secondary Hyperalgesia

Secondary hyperalgesia develops after a wide range of cutaneous injuries including burns, mechanical trauma, and freeze injuries. One technique to produce secondary hyperalgesia that is used by a number of investigators is to inject a small volume (10  $\mu\text{L}$ ) of capsaicin into the skin. Capsaicin is the active ingredient in hot peppers and produces intense pain upon injection. Once the pain disappears, a large area of flare and secondary hyperalgesia becomes apparent (Fig. 6).

The flare, or reddening of the skin, is due to a dilation of the blood vessels in the skin and lasts for about 30 min. The activation of nociceptors by capsaicin leads to the release of substances, called neuropeptides, from the terminals of the nociceptors. These neuropeptides cause vasodilation. The release of neuropeptides is not restricted to the site of capsaicin injection. It also occurs in other terminals of these nociceptors outside the injection area. Excitation of one part of the receptor by capsaicin leads to the propagation of signals to other parts of the receptor and the release of the neuropeptides. The large area of the flare, relative to the area to which capsaicin was applied, reflects the large area over which these particular nociceptors have terminals.

Two forms of secondary hyperalgesia to mechanical stimuli are observed (Fig. 6). *Stroking hyperalgesia* occurs in response to gentle stroking of the skin with soft stimuli (e.g., a Q-tip). This is sometimes called dynamic hyperalgesia because it involves a moving stimulus. *Punctate hyperalgesia* occurs in response to punctate mechanical



	Stroking hyperalgesia (dynamic)	Punctate hyperalgesia (static)
Adequate stimulus	Light stroking	Punctate stimuli
Area	Small	Large
Duration	Short	Long

**Figure 6.** Two forms of secondary hyperalgesia. (top) A tissue injury (or capsaicin injection) leads to the development of a large area over which the pain to punctate stimuli is enhanced (punctate hyperalgesia). Within this area is a smaller zone in which there is pain to gentle stroking of the skin (stroking hyperalgesia). An area of reddening or flare is also apparent, which is due to vasodilation. (bottom) The properties of punctate and stroking hyperalgesia differ, suggesting different neural mechanisms.

stimuli such as a pin or a nylon monofilament. Punctate hyperalgesia is sometimes called static hyperalgesia because it involves applying a stimulus to a fixed position for a period of time.


The characteristics of stroking and punctate hyperalgesia differ. The area of stroking hyperalgesia is smaller than the area of punctate hyperalgesia. In addition, the duration of stroking hyperalgesia is short (e.g., about 30 min after capsaicin injection), whereas the duration of punctate hyperalgesia is long (up to 24 h after capsaicin injection). This difference provides the first line of evidence that the neural code for the two forms of hyperalgesia differs.

We wondered whether peripheral sensitization accounts for the secondary hyperalgesia that develops in


uninjured skin adjacent to an injury. It could be that the sensitization that occurs at the site of injury spreads to adjacent tissue. For example, do the neuropeptides that are released in the area of flare lead to sensitization of the nociceptors? To investigate this, we (and others) applied injuries to part of the receptive area of the nociceptors under study.<sup>8,9</sup> Sensitization occurs in the injured part of the receptor, but the responses of the uninjured part are not altered (Fig. 7). Thus, sensitization of the peripheral nociceptors does not account for secondary hyperalgesia. Rather, the preponderance of evidence indicates that secondary hyperalgesia is due to changes in the central processing of nociceptive information. We call this “central sensitization.”

One of the most compelling experiments to demonstrate central sensitization is illustrated in Fig. 8.<sup>10</sup> Here, a local anesthetic was injected into the nerve serving the right forearm near the elbow. Capsaicin was then injected into the anesthetized area. There was no pain because the parent nerve was anesthetized, but a flare developed, indicating that the nociceptors had been activated (Fig. 8a). At the same time, the left arm (control) was injected with capsaicin. This produced intense pain, flare, and a large zone of secondary hyperalgesia (Fig. 8b). After the anesthetic had worn off (i.e., 3 h later), the skin was tested again. The left arm still exhibited a large zone of hyperalgesia. In the right arm, the mechanical stimuli felt normal; there was no hyperalgesia.

This experiment demonstrates that anesthetizing the peripheral nerve prevents the development of secondary hyperalgesia. We postulate that the barrage of input from nociceptors at the time of capsaicin injection leads to the development of sensitization in the central nervous system. The anesthetic blocks the signals from the nociceptors from reaching the brain. The nociceptors are still activated since a flare developed, but there is no sensitization in the peripheral tissues.



**Figure 7.** Tissue injury leads to sensitization of nociceptive fibers at the site of injury, but there is no spread of sensitization to nociceptive terminals outside of the injury site.



**Figure 8.** A nerve block prevents development of secondary hyperalgesia. (a) A local anesthetic was injected into a nerve near the elbow to produce a large zone of anesthesia in the skin (shaded area). Capsaicin was then injected into the anesthetized area and produced a flare but no pain (left panel). At 3 h later, mechanical testing of the skin revealed no hyperalgesia. (b) Capsaicin was also injected into the opposite arm in the absence of a nerve block and produced a flare and a large zone of hyperalgesia to punctate and stroking stimuli. At 3 h later, the hyperalgesia to punctate stimuli was still present. Solid lines = areas of hyperalgesia to punctate mechanical stimuli; dotted lines = areas of hyperalgesia to light touch; dashed lines = areas of flare. (Used with permission from Ref. 10.)

Signs of central sensitization are already apparent at the spinal cord, where the first synapse in the pain system occurs. Neurons at this level exhibit enhanced responsiveness to mechanical stimuli applied outside the injury area.

### Punctate Hyperalgesia Signaled by A-Fiber Nociceptors

Before considering which peripheral nerve fibers provide the signal for punctate hyperalgesia, we first


consider which nerve fibers signal the pain to punctate stimuli in normal skin. We performed psychophysical experiments in which sharp probes of different forces were applied to the back of the hand of normal volunteers.<sup>11</sup> Each probe consisted of a 200- $\mu\text{m}$ -dia. stainless steel wire attached to a rod with a given weight. The weights were varied so that the probes delivered forces ranging from 8 to 512 mN in factor of 2 increments. The probes were applied for 1 s in random order. The subjects reported that these objects produced a sharp pain sensation. They were asked to rate the magnitude of the pain using a 0–100 numerical scale where “0” corresponds to no pain and “100” corresponds to the most intense pain imaginable. The stimulus-response function is shown in Fig. 9a. The threshold for pain was around 16 mN, and the pain ratings increased monotonically with increasing force.

In the same subjects, a fixed pressure was applied to the nerve that supplies the top of the hand. After approximately 45 min, this pressure resulted in a selective block of conduction in A-fibers, but C-fibers still conducted. (This observation is similar to the phenomenon most readers probably have experienced when they wake up in the middle of the night and can’t feel anything in their hand because they have been sleeping on it.) The punctate stimuli were again applied to the hand. The pressure block resulted in a significant decrease in the pain ratings reported by the subjects (Fig. 9a). The pain threshold increased, and the response to suprathreshold stimuli decreased. Thus, when A-fibers are blocked but C-fibers are still conducting, pain ratings to punctate stimuli go down substantially. These data provide psychophysical evidence that A-fiber nociceptors perform a primary role in signaling the pain to punctate stimuli in normal skin.

How do the A-fiber and C-fiber nociceptors respond to these types of stimuli? A Laboratory-developed mechanical stimulator system (see the boxed insert, “Mechanical Stimulator for Studies of Pain”) was employed to apply controlled force stimuli to the skin area where the A-fiber and C-fiber nociceptors terminated.<sup>12</sup> The average response to punctate mechanical stimuli for both nociceptors is shown in Fig. 9b.<sup>13</sup> In these experiments, a 400- $\mu\text{m}$  cylindrical probe was applied to the skin for a period of 3 s, and the stimuli were applied as an ascending series from 40 to 200 mN in 40-mN increments. The response of the A-fiber and C-fiber nociceptors to these forces is considerably different. The response of the A-fibers increases monotonically over this stimulus range, much like the pain ratings did in Fig. 9a. In contrast, the response of the C-fibers reaches a plateau at the higher forces, although pain sensation keeps increasing in that range. These data provide electrophysiological evidence that A-fibers play a major role in signaling the sharp pain to punctate stimuli in normal skin.

#### *Punctate Hyperalgesia Signaled by a Subclass of A-Fiber Nociceptors*

We now consider whether A- or C-fiber nociceptors signal the enhanced pain to punctate stimuli that develops in the zone of secondary hyperalgesia. To address this question, we made use of another property of capsaicin. Topical application of capsaicin can lead to a desensitization of the skin (desensitization also explains why people who eat hot foods frequently can tolerate higher doses of hot peppers than those who don’t). A 10% capsaicin cream was applied under an occlusive dressing to a small area of the



**Figure 9.** A-fiber nociceptors signal sharp pain to punctate stimuli in normal skin. (a) Pain ratings of human subjects to punctate mechanical stimuli. The threshold for pain was around 16 mN, and the pain ratings increased monotonically with increasing force (red curve). When nerve conduction in the A-fibers was blocked, the pain ratings decreased substantially (purple curve). (b) Responses of nociceptors in monkeys to punctate mechanical stimuli. The responses of A-fiber nociceptors increased monotonically with stimulus force. The responses of C-fiber nociceptors reached a plateau at higher stimulus forces.




### MECHANICAL STIMULATOR FOR STUDIES OF PAIN

A computer-based electromechanical stimulator system was developed at APL for our neurophysiological and psychophysical studies of pain.<sup>12</sup> The core of the stimulator is a servo-controlled linear motor capable of generating 1 kg of force over a 22-mm range. Three load cells (resolution = 1/8 g, range = 250 g) arranged in an equilateral triangle are attached to one end of the shaft. The interchangeable probe tips are attached at the center of the triangle. Forces collinear and tangential to the probe are calculated using the signal from these three sensors. An optical encoder (resolution = 1  $\mu$ m, range = 25 mm) is positioned on the other end of the shaft and used to measure probe position. A microprocessor-based digital control system permits smooth switching of feedback control between force or position at the 1-kHz update rate.

The stimulator is mounted on a microprocessor-controlled three-axis translation system. This system is used to automatically move the probe to specific locations on the skin over a range of greater than 15 cm to an accuracy of better than 10  $\mu$ m. The stimulator can be programmed to move in a coordinate system parallel to the skin surface being examined.

Photograph of the mechanical stimulator. (Reprinted from Ref. 12 with permission, © 1995, Elsevier.)




forearm (25 × 25 mm) for 6 h on 2 consecutive days (Fig. 10a). A second area was treated identically with a control cream that did not contain capsaicin. The two areas were separated by a 10-mm strip of untreated skin.

As shown in Fig. 10b, the capsaicin treatment produced a marked desensitization of the skin to heat stimuli. Pain ratings to an intense heat stimulus (53°C, 4 s) were almost abolished after 2 days of treatment. Pain ratings to heat remained low for more than 2 weeks. Others have shown that topical capsaicin treatment leads to a degeneration of nerve fiber terminals in the skin,<sup>14</sup> indicating that this treatment results in a long-lasting desensitization of AMH and CMH nociceptors.

One day after the pretreatment was completed, capsaicin (40  $\mu$ g) was injected into the untreated area between the two treatment sites. This capsaicin injection led to the development of a large zone of secondary hyperalgesia as well as a large area of flare. The hyperalgesia extended symmetrically around the injection site and included the vehicle and capsaicin pretreatment sites. In contrast, the flare extended into the vehicle but not into the capsaicin treatment area. Although capsaicin treatment caused a desensitization of mechano-heat-sensitive nociceptors, as evidenced by the substantial reduction in heat pain ratings and the absence of a flare response, secondary hyperalgesia was not affected.

In another study using capsaicin to desensitize an area on the forearm,<sup>15</sup> injection of capsaicin between the capsaicin pretreatment area and the vehicle pretreatment area again led to a large zone of secondary hyperalgesia that encompassed the capsaicin pretreatment area. In this study, we obtained pain ratings in response to the application of a blade-shaped probe to the skin with our mechanical stimulator system. Pain ratings increased dramatically after injection of the capsaicin, indicating the presence of mechanical hyperalgesia. The pain ratings at the capsaicin pretreatment area were not significantly different from those at the vehicle pretreatment area (Figs. 10c and 10d). Based on these results, we conclude that hyperalgesia persists in capsaicin-treated areas and that mechano-heat-sensitive nociceptors do not signal punctate hyperalgesia.

In subsequent experiments, we applied a pressure block to the nerve to obstruct A-fiber conduction. We found that hyperalgesia to punctate stimuli disappeared when A-fiber conduction was blocked, but C-fibers were still conducting.<sup>16</sup> We conclude from these experiments that A-fiber nociceptors signal punctate hyperalgesia. These A-fibers must be a subset of the A-fiber nociceptors that are neither capsaicin sensitive nor heat sensitive. Electrophysiological experiments are under way to characterize the properties of these fibers.



**Figure 10.** Hyperalgesia to punctate stimuli persists in capsaicin-desensitized skin. (a) Capsaicin cream was applied to a skin area over 2 days to desensitize the skin. A second area was treated with a vehicle. Capsaicin was then injected between the two areas. Secondary hyperalgesia to punctate stimuli developed symmetrically around the capsaicin injection site. The flare extended into the vehicle-treated area but not the capsaicin-treated area. (b) Heat testing revealed that the topical capsaicin treatment led to a decreased sensitivity to heat stimuli. (c) Pain ratings to punctate stimuli applied to the vehicle-treated site increased dramatically after capsaicin injection. (d) Pain ratings applied to the capsaicin-treated site also increased dramatically after capsaicin injection. (Adapted from Ref. 11 with permission, © 2001, Oxford University Press.)

## CONCLUSIONS

Nerve fiber terminals in the skin that respond selectively to noxious stimuli are called nociceptors. Nociceptors are thought to be responsible for signaling pain sensation. Two classes of nociceptors can be identified based on the speed of action potential propagation in the parent nerve fiber: A-fiber nociceptors (fast) and C-fiber nociceptors (slow). A-fiber nociceptors are responsible for sharp pain sensation in response to punctate mechanical stimuli. C-fiber nociceptors are responsible for the burning pain sensation from hot stimuli.

Enhanced pain or hyperalgesia occurs after injury to the skin. Primary hyperalgesia develops at the site of injury and is characterized by enhanced pain to mechanical and heat stimuli. Secondary hyperalgesia occurs in

uninjured skin surrounding the site of injury and is characterized by enhanced pain to mechanical, but not heat, stimuli. The neural mechanisms of primary and secondary hyperalgesia differ.

An injury (e.g., a burn) to a nociceptor leads to an enhanced responsiveness of the nociceptor called sensitization. This sensitization occurs only for those parts of the nociceptor that are directly injured. Thus, sensitization of nociceptors is thought to account for primary hyperalgesia. Nociceptors in the area of secondary hyperalgesia are not sensitized. Secondary hyperalgesia is thought to be a result of sensitization in the central nervous system.

Hyperalgesia is prevalent in many disease states. The hyperalgesia that occurs in inflammation (e.g., rheumatoid arthritis) is comparable to primary hyperalgesia and is thought to result from nociceptor sensitization. The hyperalgesia to light touch that occurs in neuropathic pain (e.g., shingles) is comparable to secondary hyperalgesia and is thought to be the result of central sensitization.

**ACKNOWLEDGMENTS:** Several instruments used in this research were developed at APL. The initial concepts for the laser stimulator were developed by Drs. Ronald Walker and Vernon Mountcastle. The mechanical stimulator system was developed by Wolfgang Schneider. We also thank the many postdoctoral fellows and research assistants who have worked on various aspects of this research, including Zahid Ali, Karen Davis, Perry Fuchs, Timothy Hartke, Sylvia Horasek, Jason Huang, Sherif Meleka, Yuan Bo Peng, Carolina Roza, Robert Slugg, Donna Bea Tillman, Brian Turnquist, and Gang Wu. Some of these studies were done in collaboration with Rolf-Detlef Treede and Walter Magerl (University of Mainz, Germany). This research was supported by NIH.


## REFERENCES

- Ali, Z., Meyer, R. A., and Campbell, J. N., "Secondary Hyperalgesia to Mechanical but Not Heat Stimuli Following a Capsaicin Injection in Hairy Skin," *Pain* 68(2-3), 401-411 (1996).
- Raja, S. N., Campbell, J. N., and Meyer, R. A., "Evidence for Different Mechanisms of Primary and Secondary Hyperalgesia Following Heat Injury to the Glabrous Skin," *Brain* 107, 1179-1188 (1984).
- Meyer, R. A., and Campbell, J. N., "Myelinated Nociceptive Afferents Account for the Hyperalgesia That Follows a Burn to the Hand," *Science* 213, 1527-1529 (1981).
- Meyer, R. A., Walker, R. E., and Mountcastle, V. B., "A Laser Stimulator for the Study of Cutaneous Thermal Pain Sensation," *IEEE Trans. Biomed. Eng.* 23, 54-60 (1976).
- Davis, K. D., Meyer, R. A., Turnquist, J. L., Filloon, T. G., Pappagallo, M., and Campbell, J. N., "Cutaneous Injection of the Capsaicin Analog, NE-21610, Produces Analgesia to Heat but Not to Mechanical Stimuli in Man," *Pain* 62, 17-26 (1995).
- Meyer, R. A., and Campbell, J. N., "Peripheral Neural Coding of Pain Sensation," *Johns Hopkins APL Tech. Dig.* 2, 164-171 (1981).

- <sup>7</sup>Peng, Y. B., Ringkamp, M., Meyer, R. A., and Campbell, J. N., "Fatigue and Paradoxical Enhancement of Heat Response in C-Fiber Nociceptors from Cross-Modal Excitation," *J. Neurosci.* **23**(11), 4766–4774 (2003).
- <sup>8</sup>Campbell, J. N., Khan, A. A., Meyer, R. A., and Raja, S. N., "Responses to Heat of C-Fiber Nociceptors in Monkey Are Altered by Injury in the Receptive Field but Not by Adjacent Injury," *Pain* **32**, 327–332 (1988).
- <sup>9</sup>Schmelz, M., Schmidt, R., Ringkamp, M., Forster, C., Handwerker, H. O., and Torebjörk, H. E., "Limitation of Sensitization to Injured Parts of Receptive Fields in Human Skin C-Nociceptors," *Exp. Brain Res.* **109**(1), 141–147 (1996).
- <sup>10</sup>LaMotte, R. H., Shain, C. N., Simone, D. A., and Tsai, E.-F.P., "Neurogenic Hyperalgesia: Psychophysical Studies of Underlying Mechanisms," *J. Neurophysiol.* **66**, 190–211 (1991).
- <sup>11</sup>Magerl, W., Fuchs, P. N., Meyer, R. A., and Treede, R-D., "Roles of Capsaicin-Insensitive Nociceptors in Cutaneous Pain and Secondary Hyperalgesia," *Brain* **124**(Pt. 9), 1754–1764 (2001).
- <sup>12</sup>Schneider, W., Slugg, R. M., Turnquist, B. P., Meyer, R. A., and Campbell, J. N., "An Electromechanical Stimulator System for Neurophysiological and Psychophysical Studies of Pain," *J. Neurosci. Methods* **60**, 61–68 (1995).
- <sup>13</sup>Slugg, R. M., Meyer, R. A., and Campbell, J. N., "Response of Cutaneous A- and C-Fiber Nociceptors in the Monkey to Controlled-Force Stimuli," *J. Neurophysiol.* **83**(4), 2179–2191 (2000).
- <sup>14</sup>Nolano, M., Simone, D. A., Wendelschafer-Crabb, G., Johnson, T., Hazen, E., and Kennedy, W. R., "Topical Capsaicin in Humans: Parallel Loss of Epidermal Nerve Fibers and Pain Sensation," *Pain* **81**(1-2), 135–145 (1999).
- <sup>15</sup>Fuchs, P. N., Campbell, J. N., and Meyer, R. A., "Secondary Hyperalgesia Persists in Capsaicin Desensitized Skin," *Pain* **84**(2-3), 141–149 (2000).
- <sup>16</sup>Ziegler, E. A., Magerl, W., Meyer, R. A., and Treede, R-D., "Secondary Hyperalgesia to Punctate Mechanical Stimuli: Central Sensitization to A-Fibre Nociceptor Input," *Brain* **122**(Pt. 12), 2245–2257 (1999).

## THE AUTHORS

**Richard A. Meyer** is a member of the Principal Professional Staff in APL's National Security Technology Department. He is a Professor in the Neurological Surgery and Biomedical Engineering departments at the Johns Hopkins Medical School and Chairman of the school's Blaustein Pain Treatment Center Research Committee. His primary research interest is in the peripheral neural mechanisms of pain sensation. **Matthias Ringkamp** is an Assistant Professor in the Department of Neurological Surgery at the Medical School. Dr. Ringkamp's research interest is in the neural mechanisms of pain and itch sensation. **James N. Campbell** serves as Professor and Vice Chair in the Department of Neurological Surgery at the Medical School. Dr. Campbell is the Director of the Blaustein Pain Treatment Center and President of the American Pain Foundation. His clinical practice specializes in spine, spinal cord, and peripheral nerve surgery. Dr. Campbell's area of research is the neurobiology of pain. **Srinivasa N. Raja** is a Professor of Anesthesiology at the Hopkins Medical School. Dr. Raja is also the Director of the Pain Division in the Department of Anesthesiology. His clinical interests include regional anesthesia and the management of acute and chronic pain states (e.g., sympathetically maintained pain, postherpetic neuralgia, post-amputation pain). His research efforts are aimed at understanding the peripheral and central mechanisms of neuropathic pain. Further information on pain research activities at APL can be obtained from Richard A. Meyer. His e-mail address is richard.meyer@jhuapl.edu.



From left: Matthias Ringkamp, Srinivasa N. Raja, Richard A. Meyer, and James N. Campbell.